

From N -Body Schrödinger to Vlasov

François Golse

CMLS, École polytechnique & CNRS, Université Paris-Saclay

Séminaire "EDP et applications"
Collège de France, June 3rd 2016

Work with T. Paul, [arXiv:1510.06681](https://arxiv.org/abs/1510.06681)

The Vlasov equation with Lipschitz continuous interaction force has been derived from the N -body problem of classical mechanics in the large N , small coupling constant limit (Neunzert-Wick 1973, Braun-Hepp 1977, Dobrushin 1979)


Problem: Is it possible to derive the Vlasov equation from the quantum N -body problem by a joint semiclassical ($\hbar \rightarrow 0$) and mean field ($N \rightarrow \infty$) limit?

[Graffi-Martinez-Pulvirenti M3AS 2003]

[Pezzotti-Pulvirenti Ann IHP 2009]

[Benedikter-Porta-Saffirio-Schlein arXiv:1502.04230]

The diagram



Uniformity as $\hbar \rightarrow 0$ of the upper horizontal (mean-field) limit:
[FG-Mouhot-Paul CMP **343** (2016), 165–205]

QUANTUM VS CLASSICAL DYNAMICS

Quantum vs classical dynamics

Heisenberg/Von Neumann equation

$$i\hbar\partial_t\rho = [\mathcal{H}, \rho]$$

where $\mathcal{H} = \mathcal{H}^*$ = Hamiltonian, while $[A, B] := AB - BA$ and

$$\rho = \rho^* \geq 0, \quad \text{Tr}_{\mathfrak{H}} \rho = 1 \Leftrightarrow \rho \in \mathcal{D}(\mathfrak{H}) \text{ with } \mathfrak{H} := L^2(\mathbf{R}^d)$$

Liouville equation

$$\partial_t f + \{H, f\} = 0 \quad H \equiv H(x, \xi) \in \mathbf{R}$$

$$f \equiv f(t, x, \xi) \geq 0, \quad \iint_{\mathbf{R}^d \times \mathbf{R}^d} f(t, x, \xi) dx d\xi = 1$$

with

$$\{\mathcal{H}, f\} := \nabla_{\xi} H \cdot \nabla_x f - \nabla_x H \cdot \nabla_{\xi} f$$

Comparing quantum and classical densities

- WKB ansatz, superpositions of coherent states driven by the classical dynamics produce (local in the case of WKB) **approximate** solutions of the Schrödinger equation, i.e. with source terms
- Wigner transform of $\rho \in \mathcal{D}(\mathfrak{H})$

$$W_{\hbar}[\rho](x, \xi) := \frac{1}{(2\pi)^d} \int_{\mathbf{R}^d} e^{-i\xi \cdot y} \rho(x + \frac{1}{2}\hbar y, x - \frac{1}{2}\hbar y) dy$$

satisfies a transport-like equation but is in general not nonnegative

- Husimi transform

$$\tilde{W}_{\hbar}[\rho] := e^{\hbar\Delta_{x,\xi}/4} W_{\hbar}[\rho] \geq 0$$

satisfies a transport-like equation involving the complex extension of the potential

Coupling quantum and classical densities

Following Dobrushin's 1979 derivation of Vlasov's equation, seek to measure the difference between the quantum and the classical dynamics by a Monge-Kantorovich type distance.

Couplings of $\rho \in \mathcal{D}(\mathfrak{H})$ and p probability density on $\mathbf{R}^d \times \mathbf{R}^d$

$$(x, \xi) \mapsto Q(x, \xi) = Q(x, \xi)^* \in \mathcal{L}(\mathfrak{H}) \text{ s.t. } Q(x, \xi) \geq 0$$

$$\text{Tr}(Q(x, \xi)) = p(x, \xi), \quad \iint_{\mathbf{R}^d \times \mathbf{R}^d} Q(x, \xi) dx d\xi = \rho$$

The set of all couplings of the densities ρ and p is denoted $\mathcal{C}(\rho, p)$

Pseudo-distance between quantum and classical densities

Cost function comparing classical and quantum “coordinates” (i.e. position and momentum)

$$c_{\hbar}(x, \xi) := |x - y|^2 + |\xi + i\hbar\nabla_y|^2$$

Define a pseudo-distance “à la” Monge-Kantorovich

$$E_{\hbar}(p, \rho) := \left(\inf_{Q \in \mathcal{C}(p, \rho)} \iint_{\mathbf{R}^d \times \mathbf{R}^d} \text{Tr}(c_{\hbar}(x, \xi) Q(x, \xi)) dx d\xi \right)^{1/2}$$

Monge-Kantorovich(-Rubinshtein) or Vasershtein distances

Let $p \geq 1$ and $\mu, \nu \in \mathcal{P}_p(\mathbf{R}^d)$ with bounded moment of order p

Coupling of μ, ν : any $\pi \in \mathcal{P}(\mathbf{R}^d \times \mathbf{R}^d)$ s.t.

$$\iint (\phi(x) + \psi(y))\pi(dxdy) = \int \phi(x)\mu(dx) + \int \psi(y)\nu(dy)$$

Set of couplings of μ, ν denoted $\Pi(\mu, \nu)$; define

$$\text{dist}_{MK,p}(\mu, \nu) = \inf_{\pi \in \Pi(\mu, \nu)} \left(\iint |x - y|^p \pi(dxdy) \right)^{1/p}$$

This distance metrizes the topology of weak convergence on $\mathcal{P}_p(\mathbf{R}^d)$

- Coherent state with $q, p \in \mathbf{R}^d$:

$$|q + ip, \hbar\rangle = (\pi\hbar)^{-d/4} e^{-|x-q|^2/2\hbar} e^{ip \cdot x/\hbar}$$

- With the identification $z = q + ip \in \mathbf{C}^d$

$$\text{OP}^T(\mu) := \frac{1}{(2\pi\hbar)^d} \int_{\mathbf{C}^d} |z, \hbar\rangle \langle z, \hbar| \mu(dz), \quad \text{OP}^T(1) = I$$

- Fundamental properties:

$$\mu \geq 0 \Rightarrow \text{OP}^T(\mu) \geq 0, \quad \text{Tr}(\text{OP}^T(\mu)) = \frac{1}{(2\pi\hbar)^d} \int_{\mathbf{C}^d} \mu(dz)$$

- Important formulas:

$$W_{\hbar}[\text{OP}^T(\mu)] = \frac{1}{(2\pi\hbar)^d} e^{\hbar\Delta_{q,p}/4} \mu, \quad \tilde{W}_{\hbar}[\text{OP}^T(\mu)] = \frac{1}{(2\pi\hbar)^d} e^{\hbar\Delta_{q,p}/2} \mu$$

Basic properties of the pseudo-distance E_{\hbar}

Thm A Let $\rho =$ probability density on $\mathbf{R}^d \times \mathbf{R}^d$ s.t.

$$\iint_{\mathbf{R}^d \times \mathbf{R}^d} (|x|^2 + |\xi|^2) \rho(x, \xi) dx d\xi < \infty$$

(1) For each $\rho \in \mathcal{D}(\mathfrak{H})$ one has $E_{\hbar}(\rho, \rho) \geq \frac{1}{2} d \hbar$

(2) For each $\mu \in \mathcal{P}(\mathbf{R}^d \times \mathbf{R}^d)$ one has

$$E_{\hbar}(\rho, \text{OP}_{\hbar}^T((2\pi\hbar)^d \mu))^2 \leq \text{dist}_{\text{MK},2}(\rho, \mu)^2 + \frac{1}{2} d \hbar$$

(3) For each $\rho \in \mathcal{D}(\mathfrak{H})$, one has

$$E_{\hbar}(\rho, \rho)^2 \geq \text{dist}_{\text{MK},2}(\rho, \tilde{W}_{\hbar}[\rho])^2 - \frac{1}{2} d \hbar$$

(4) If $\rho_{\hbar} \in \mathcal{D}(\mathfrak{H})$ and $W_{\hbar}[\rho_{\hbar}] \rightarrow \mu$ in \mathcal{S}' , then $\mu \in \mathcal{P}(\mathbf{R}^d \times \mathbf{R}^d)$ and

$$\liminf_{\hbar \rightarrow 0} E_{\hbar}(\rho, \rho) \geq \text{dist}_{\text{MK},2}(\rho, \mu)$$

QUANTITATIVE CONVERGENCE RATE

Hartree vs Vlasov equations

Hartree equation for density matrices

$$\partial_t \rho_{\hbar}(t) = -\frac{i}{\hbar} [\mathcal{H}[\rho_{\hbar}(t)], \rho_{\hbar}(t)]$$

with

$$\mathcal{H}[\rho] = -\frac{1}{2} \hbar^2 \Delta + \int_{\mathbf{R}^d} V(x-z) \rho(z, z) dz$$

Vlasov equation for $f \equiv f(t, x, \xi)$ probability density

$$\partial_t f = -\{H_f, f\} = -\xi \cdot \nabla_x f + \nabla_x V_f \cdot \nabla_{\xi} f$$

with

$$V_f(t, x) = \iint_{\mathbf{R}^d} V(x-z) \rho[f](t, z) dz, \quad \text{where } \rho[f] := \int_{\mathbf{R}^d} f d\xi$$

Thm B Let V be an even, real-valued function of class $C^{1,1}$ on \mathbf{R}^d . Let $f^{in} \equiv f^{in}(x, \xi) \in L^1(|x|^2 + |\xi|^2) dx d\xi$ be a probability density on $\mathbf{R}^d \times \mathbf{R}^d$, and let f be the solution of the Vlasov equation with initial data f^{in} . Let ρ_{\hbar} be a solution of the Hartree equation with initial data $\rho_{\hbar}^{in} \in \mathcal{D}(\mathfrak{H})$. Then

$$E_{\hbar}(f(t), \rho_{\hbar}(t))^2 \leq e^{\Lambda t} E_{\hbar}(f^{in}, \rho_{\hbar}^{in})^2$$

with $\Lambda := 1 + 4 \max(1, \text{Lip}(\nabla V))$. If $\rho_{\hbar}^{in} = \text{OP}_{\hbar}^T((2\pi\hbar)^d \mu^{in})$, then

$$\text{dist}_{\text{MK},2}(f(t), \tilde{W}_{\hbar}[\rho_{\hbar}(t)])^2 \leq e^{\Lambda t} (\text{dist}_{\text{MK},2}(f^{in}, \mu^{in})^2 + \frac{1}{2} d\hbar) + \frac{1}{2} d\hbar$$

[P.-L. Lions-T. Paul, Rev. Mat. Iberoam. 1993]: non quantitative treatment including Coulomb

N -body von Neumann and Liouville equations

N -body von Neumann equation

$$\partial_t \rho_{N,\hbar} = -\frac{i}{\hbar} [\mathcal{H}_N, \rho_{N,\hbar}]$$

where $\rho_{N,\hbar} \in \mathcal{D}(\mathfrak{H}_N)$, with $\mathfrak{H}_N = \mathfrak{H}^{\otimes N} = L^2((\mathbf{R}^d)^N)$ and

$$\mathcal{H}_N := \sum_{j=1}^N -\frac{1}{2}\hbar^2 \Delta_{y_j} + \frac{1}{N} \sum_{1 \leq j < k \leq N} V(y_j - y_k)$$

N -body Liouville equation

$$\partial_t f_N = -\{H_N, f_N\} = -\sum_{j=1}^N \xi_j \cdot \nabla_{x_j} f_N + \frac{1}{N} \sum_{j,k=1}^N \nabla V(x_j - x_k) \cdot \nabla_{\xi_j} f_N$$

Indistinguishable particles and symmetries

Notation

$$\begin{aligned} X_N &:= (x_1, \dots, x_N), & \Xi_N &:= (\xi_1, \dots, \xi_N) \\ \sigma \cdot X_N &:= (x_{\sigma(1)}, \dots, x_{\sigma(N)}) & \text{for } \sigma \in \mathfrak{S}_N \end{aligned}$$

Classical N -body symmetric probability density: for all $t \geq 0$

$$f_N(t, \sigma \cdot X_N, \sigma \cdot \Xi_N) = f_N(t, X_N, \Xi_N) \quad \text{for all } \sigma \in \mathfrak{S}_N$$

Quantum symmetric N -body density: for all $t \geq 0$

$$U_\sigma \rho_{N, \hbar}(t) U_\sigma^* = \rho_{N, \hbar}(t)$$

where U_σ is the operator on \mathfrak{H}_N defined by

$$U_\sigma \psi(X_N) = \psi(\sigma \cdot X_N)$$

Symmetric densities and k -particle marginals

For $\rho_N \in \mathcal{D}^s(\mathfrak{H}_N)$, its k -particle marginal is $\rho_N^k \in \mathcal{D}^s(\mathfrak{H}_k)$ such that

$$\mathrm{Tr}_{\mathfrak{H}_k}(A\rho_N^k) = \mathrm{Tr}_{\mathfrak{H}_N}((A \otimes I_{\mathfrak{H}_{N-k}})\rho_N)$$

for each $A \in \mathcal{L}(\mathfrak{H}_k)$

Symmetric classical density $f_N \equiv f_N(X_N, \Xi_N)$; its k -particle marginal:

$$f_N^k(X_k, \Xi_k) = \int f_N(X_N, \Xi_N) dx_{k+1} d\xi_{k+1} \dots dx_N d\xi_N$$

From von Neumann to Liouville uniformly in N

Thm C Let V be an even, real-valued function of class $C^{1,1}$ on \mathbf{R}^d . Let $\rho_{N,\hbar}^{in} \in \mathcal{D}^s(\mathfrak{H}_N)$ and F_N^{in} be a symmetric probability density on $(\mathbf{R}^d \times \mathbf{R}^d)^N$ in $L^1(|X_N|^2 + |\Xi_N|^2)dX_Nd\Xi_N)$. Let $\rho_{N,\hbar}$ and F_N be the solutions of the von Neumann and the Liouville equations resp. with initial data $\rho_{N,\hbar}^{in}$ and F_N^{in} .

Setting $\Lambda = 1 + 4 \max(1, \text{Lip}(\nabla(V)))^2$, one has

$$E_{\hbar}(F^1(t), \rho_{N,\hbar}^1(t))^2 \leq \frac{1}{N} E_{\hbar}(F_N^{in}, \rho_{N,\hbar}^{in})^2 e^{\Lambda t}$$

If $\rho_{N,\hbar}^{in} = \text{OP}_{\hbar}^T[(2\pi\hbar)^{Nd} \mu^{in}]$, then

$$\begin{aligned} & \text{dist}_{\text{MK},2}(F^1(t), \tilde{W}_{\hbar}[\rho_{N,\hbar}^1(t)])^2 \\ & \leq \left(\frac{1}{N} \text{dist}_{\text{MK},2}((F_N^{in}, \tilde{W}_{\hbar}[\rho_{N,\hbar}^{in}]) + \frac{1}{2}d\hbar) \right) e^{\Lambda t} + \frac{1}{2}d\hbar \end{aligned}$$

From N -body von Neumann to Vlasov

Thm D Let $f^{in} \equiv f^{in}(x, \xi) \in L^1(|x|^2 + |\xi|^2) dx d\xi$ be a probability density on $\mathbf{R}^d \times \mathbf{R}^d$, an $\rho_{N, \hbar}^{in} \in \mathcal{D}^s(\mathfrak{H}_N)$. Let f and $\rho_{N, \hbar}$ be the solutions of the Vlasov equation and the von Neumann equation resp. with initial data f^{in} and $\rho_{N, \hbar}^{in}$.

$$E_{\hbar}(f(t), \rho_{\hbar, N}^1(t))^2 \leq \frac{1}{N} E_{\hbar}((f^{in})^{\otimes N}, \rho_{\hbar, N}^{in})^2 e^{\Gamma t} + \frac{(2\|\nabla V\|_{L^\infty})^2}{N-1} \frac{e^{\Gamma t} - 1}{\Gamma}$$

If moreover $\rho_{\hbar, N}^{in} = \text{OP}_{\hbar}^T [(2\pi\hbar)^{dN} (f^{in})^{\otimes N}]$

$$\text{dist}_{\text{MK}, 2}(f(t), \widetilde{W}_{\hbar}[\rho_{\hbar, N}^1(t)])^2 \leq \frac{1}{2} d\hbar(1 + e^{\Gamma t}) + \frac{(2\|\nabla V\|_{L^\infty})^2}{N-1} \frac{e^{\Gamma t} - 1}{\Gamma}$$

Here $\Gamma = 2 + 4 \max(1, \text{Lip}(\nabla(V)))^2$.

SKETCH OF THE PROOFS

Dynamics of couplings

If F_N is a symmetric classical N -particle density and $\rho_N \in \mathcal{D}^s(\mathfrak{H}_N)$, a coupling $Q_N \in \mathcal{C}(F_N, \rho_N)$ is symmetric, denoted $Q_N \in \mathcal{C}^s(F_N, \rho_N)$

$$U_\sigma Q_N(\sigma \cdot X_N, \sigma \cdot \Xi_N) U_\sigma^* = Q_N(X_N, \Xi_N) \quad \text{for all } \sigma \in \mathfrak{S}_N$$

Let $Q_{N,\hbar}^{in} \in \mathcal{C}^s((f^{in})^{\otimes N}, \rho_{N,\hbar}^{in})$; solve

$$\partial_t Q_{N,\hbar} + \left\{ \sum_{j=1}^N H_f(x_j, \xi_j), Q_{N,\hbar} \right\} + \frac{i}{\hbar} [\mathcal{H}_N, Q_{N,\hbar}] = 0$$

with $Q_{N,\hbar}|_{t=0} = Q_{N,\hbar}^{in}$ and

$$\mathcal{H}_N := \sum_{j=1}^N -\frac{1}{2}\hbar^2 \Delta_{y_j} + \frac{1}{N} \sum_{1 \leq j < k \leq N} V(y_j - y_k)$$

$$H_f(x, \xi) := \frac{1}{2}|\xi|^2 + \iint_{\mathbb{R}^d \times \mathbb{R}^d} V(x - z) f(t, z, \zeta) dz d\zeta$$

The functional $D(t)$

Lemma For each $t \geq 0$, one has

$$Q_{N,\hbar}(t) \in \mathcal{C}^s(f(t)^{\otimes N}, \rho_{N,\hbar}(t))$$

where f is the solution of the Vlasov equation and $\rho_{N,\hbar}$ is the solution of the N -body von Neumann equation

Define

$$\begin{aligned} D(t) &:= \frac{1}{N} \iint_{(\mathbb{R}^d \times \mathbb{R}^d)^N} \sum_{k=1}^N \text{Tr}_{\mathfrak{H}^N} (c_{\hbar}(x_j, \xi_j, y_j, \nabla_{y_j}) Q_{N,\hbar}(t)) dX_N d\xi_N \\ &= \iint_{(\mathbb{R}^d \times \mathbb{R}^d)^N} \text{Tr}_{\mathfrak{H}^N} (c_{\hbar}(x_1, \xi_1, y_1, \nabla_{y_1}) Q_{N,\hbar}(t)) dX_N d\xi_N \\ &= \iint_{(\mathbb{R}^d \times \mathbb{R}^d)} \text{Tr}_{\mathfrak{H}} (c_{\hbar}(x_1, \xi_1, y_1, \nabla_{y_1}) Q_{N,\hbar}^1(t)) dx_1 d\xi_1 \\ &\geq E_{\hbar}(f(t), \rho_{N,\hbar}^1(t)) \end{aligned}$$

Multiply both sides of the equation for $Q_{N,\hbar}$ and “integrate by parts”:

$$\begin{aligned}\dot{D} &= \iint \text{Tr}_{\mathfrak{S}_1}(\{H_f(x_1, \xi_1), c_{\hbar}(x_1, \xi_1, y_1, \nabla y_1)\} Q_{N,\hbar}^1) dx_1 d\xi_1 \\ &\quad - \frac{1}{2} i \hbar \iint \text{Tr}_{\mathfrak{S}_1}([\Delta_{y_1}, c_{\hbar}(x_1, \xi_1, y_1, \nabla y_1)] Q_{N,\hbar}^1) dx_1 d\xi_1 \\ &\quad + \frac{i}{\hbar} \iint \text{Tr}_{\mathfrak{S}_2}([\frac{N-1}{N} V(y_1 - y_2), c_{\hbar}(x_1, \xi_1, y_1, \nabla y_1)]) Q_{N,\hbar}^2) dX_2 d\xi_2\end{aligned}$$

since $Q_{N,\hbar}$ is a **symmetric** coupling

In other words

$$\begin{aligned}
 \dot{D} &\leq D - \frac{N-1}{2N} \int \text{Tr}_{\mathfrak{H}_2}(Q_{N,\hbar}^2(\xi_1 + i\hbar\nabla_{y_1}) \vee \mathcal{W}(X_2, Y_2)) dX_2 d\Xi_2 \\
 &\quad - \frac{1}{2} \int \text{Tr}_{\mathfrak{H}_2}(Q_{N,\hbar}^2(\xi_1 + i\hbar\nabla_{y_1}) \vee \mathcal{V}(t, x_1, x_2)) dX_2 d\Xi_2 \\
 &\leq D - \frac{N-1}{2N} \int \text{Tr}_{\mathfrak{H}_2}(Q_{N,\hbar}^2(\xi_1 + i\hbar\nabla_{y_1}) \vee \mathcal{W}(X_2, Y_2)) dX_2 d\Xi_2 \\
 &\quad - \frac{1}{2} \int \text{Tr}_{\mathfrak{H}_2}(Q_{N,\hbar}(\xi_1 + i\hbar\nabla_{y_1}) \vee \frac{1}{N-1} \sum_{k=2}^N \mathcal{V}(t, x_1, x_k)) dX_N d\Xi_N
 \end{aligned}$$

with $a \vee b := ab + ba$ is the anticommutator while

$$\begin{aligned}
 \mathcal{V}(t, x_1, x_2) &:= \nabla V \star \rho_f(t, x_1) - \frac{N-1}{N} \nabla V(x_1, x_2) \\
 \mathcal{W}(X_2, Y_2) &:= \nabla V(x_1 - x_2) - \nabla V(y_1 - y_2)
 \end{aligned}$$

$$\begin{aligned} \dot{D} &\leq 3D + \frac{N-1}{2N} \int \text{Tr}_{\mathfrak{H}_2} (Q_{N,\hbar}^2 |\mathcal{W}(X_2, Y_2)|^2) dX_2 d\Xi_2 \\ &\quad + \frac{1}{2} \int \left| \frac{1}{N-1} \sum_{k=2}^N \mathcal{V}(t, x_1, x_k) \right|^2 \rho_f^{\otimes N} dX_N \\ &\leq 3D + \frac{N-1}{2N} L^2 \int \text{Tr}_{\mathfrak{H}_2} (Q_{N,\hbar}^2 |X_2 - Y_2|^2) dX_2 d\Xi_2 \\ &\quad + \frac{1}{2} \int \left| \frac{1}{N-1} \sum_{k=2}^N \mathcal{V}(t, x_1, x_k) \right|^2 \rho_f^{\otimes N} dX_N \\ &\leq (3 + 2L^2)D + \frac{1}{2} \int \left| \frac{1}{N-1} \sum_{k=2}^N \mathcal{V}(t, x_1, x_k) \right|^2 \rho_f^{\otimes N} dX_N \end{aligned}$$

FROM N-BODY SCHRÖDINGER TO HARTREE: UNIFORM CONVERGENCE RATE

Work with C. Mouhot & T. Paul
Commun. Math. Phys. **343** (2016), 165–205

Quantum couplings and pseudo-distance

- Density operators on a Hilbert space \mathfrak{H} :

$$\rho \in \mathcal{D}(\mathfrak{H}) \Leftrightarrow \rho = \rho^* \geq 0, \quad \text{Tr}(\rho) = 1$$

- Couplings between two density operators $\rho_1, \rho_2 \in \mathcal{D}(\mathfrak{H})$:

$$\rho \in \mathcal{D}(\mathfrak{H} \otimes \mathfrak{H}) \text{ s.t. } \begin{cases} \text{Tr}_{\mathfrak{H} \otimes \mathfrak{H}}((A \otimes I)\rho) = \text{Tr}_{\mathfrak{H}}(A\rho_1) \\ \text{Tr}_{\mathfrak{H} \otimes \mathfrak{H}}((I \otimes A)\rho) = \text{Tr}_{\mathfrak{H}}(A\rho_2) \end{cases}$$

for all $A \in \mathcal{L}(\mathfrak{H})$; the set of all such ρ will be denoted $\mathcal{Q}(\rho_1, \rho_2)$

- For $\rho_1, \rho_2 \in \mathcal{D}(L^2(\mathbf{R}^d))$, define

$$MK_2^{\hbar}(\rho_1, \rho_2) = \inf_{\rho \in \mathcal{Q}(\rho_1, \rho_2)} \text{Tr} \left(\sum_{j=1}^d ((x_j - y_j)^2 - \hbar^2 (\partial_{x_j} - \partial_{y_j})^2) \rho \right)^{1/2}$$

Dynamics of quantum couplings

Let $R_N^{in} \in \mathcal{Q}((\rho^{in})^{\otimes N}, \rho_N^{in})$ and let $t \mapsto R_N(t)$ be the solution of

$$i\hbar\partial_t R_N = [\mathbf{H}_{\rho(t)} \otimes I + I \otimes \mathcal{H}_N, R_N], \quad R_N|_{t=0} = R_N^{in}$$

Then $R_N(t) \in \mathcal{Q}((\rho(t))^{\otimes N}, \rho_N(t))$ for each $t \geq 0$. Define

$$D_N(t) = \text{Tr} \left(\frac{1}{N} \sum_{j=1}^N (Q_j^* Q_j + P_j^* P_j) R_N(t) \right)$$

with

$$Q_j = x_j - y_j, \quad P_j := \frac{\hbar}{i} (\nabla_{x_j} - \nabla_{y_j}), \quad P_j^* := \frac{\hbar}{i} (\text{div}_{x_j} - \text{div}_{y_j})$$

Theorem E

Assume that the potential V is even and satisfies $\nabla V \in W^{1,\infty}(\mathbf{R}^d)$.

Let $\rho_{\hbar}(t)$ be the solution of Hartree's equation with initial data ρ_{\hbar}^{in} , and let $\rho_{N,\hbar}(t)$ be the solution of von Neumann's equation with initial data $\rho_{N,\hbar}^{in}$ satisfying the symmetry $\rho_{N,\hbar}^{in} = U_{\sigma}^* \rho_{N,\hbar}^{in} U_{\sigma}$ for all $\sigma \in \mathfrak{S}_N$.

Then, for each $n = 1, \dots, N$, and each $t \geq 0$

$$\begin{aligned} \frac{1}{n} MK_2^{\hbar}(\rho_{\hbar}(t)^{\otimes n}, \rho_{N,\hbar}^n(t))^2 &\leq \frac{1}{N} MK_2^{\hbar}((\rho_{\hbar}^{in})^{\otimes N}, \rho_{N,\hbar}^{in})^2 e^{Lt} \\ &\quad + \frac{8}{N} \|\nabla V\|_{L^{\infty}}^2 \frac{e^{Lt} - 1}{L} \end{aligned}$$

with

$$L := 3 + 4 \operatorname{Lip}(\nabla V)^2$$

Theorem E'

Under the same assumptions as in Theorem B, assume that ρ_{\hbar}^{in} and $\rho_{N,\hbar}^{in}$ are Töplitz operators, with symbols $(2\pi\hbar)^d \mu_{\hbar}^{in}$ and $(2\pi\hbar)^{dN} \mu_{N,\hbar}^{in}$

Then, for each $n = 1, \dots, N$, and each $t \geq 0$

$$\begin{aligned} \frac{1}{n} \text{dist}_{\text{MK},2}(\tilde{W}_{\hbar}[\rho_{\hbar}(t)^{\otimes n}], \tilde{W}_{\hbar}[\rho_{N,\hbar}^n(t)])^2 &\leq \frac{8}{N} \|\nabla V\|_{L^\infty}^2 \frac{e^{Lt} - 1}{L} \\ &\quad + \frac{1}{N} \text{dist}_{\text{MK},2}((\mu_{\hbar}^{in})^{\otimes N}, \mu_{N,\hbar}^{in})^2 e^{Lt} + 2d\hbar(e^{Lt} + 1) \end{aligned}$$

with

$$L := 3 + 4 \text{Lip}(\nabla V)^2$$

- The **stability** part of the analysis (leading to the exponential amplification by Gronwall's inequality) is seen at the level of the **1st equation in the BBGKY hierarchy**
- The **consistency** part of the analysis requires distributing the interaction term \mathcal{V} on **all** the particles, and because the \mathcal{V} term depends on the X_N variables only, and the X_N marginal of $Q_{N,\hbar}$ is the N -fold tensor power of the Vlasov solution, **one concludes by (a trivial quantitative variant of the) LLN**
- Because the cost function in D is a **sum** of quantities depending on x_j, y_j, ξ_j , there is a **"localization in degree"** effect in the BBGKY hierarchy: **no Cauchy-Kovalevskaja effect when estimating D**