



PRESCRIBING THE SPECTRA
OF CUBIC GRAPHS

PETER SARNAK

LECTURE COLLEGE DE FRANCE

NOV 2023

JOINT WORK WITH

ALICIA KOLLAR AND FAN WEI

THE CONTEXT

- X A LARGE (COMPACT) LOCALLY UNIFORM GEOMETRY, \mathbb{P} A DIFFERENTIAL OPERATOR ON X (EG LAPLACIAN); WHAT GAPS IN THE SPECTRUM OF \mathbb{P} CAN BE ACHIEVED?

EXAMPLES:

- X IS LOCALLY EUCLIDEAN (GEOMETRY OF NUMBERS)
- X IS LOCALLY HYPERBOLIC
HYPERBOLIC SURFACE
HYPERBOLIC 3-MANIFOLDS
- X IS A REGULAR GRAPH
OR A BRUHAT-TITS BUILDING

WE FIXATE ON 3-REGULAR GRAPHS


• "CUBIC" GRAPHS.

CUBIC

3-REGULAR DENOTES THE SET OF ALL CONNECTED GRAPHS.

PLANAR

THE SUBSET OF CUBIC WHICH ARE PLANAR.




$\lambda_1 = -1$



$\lambda_1 = 0$


⋮

PETERSON :



$\lambda_1 = 1$

T_3 3-REGULAR TREE (UNIVERSAL COVER)



$X \in \text{CUBIC}$, $V(X)$ ⁽³⁾ VERTICES $|V(X)| = n$

ADJACENCY OPERATOR A :

$$A: \ell^2(V(X)) \rightarrow \ell^2(V(X))$$

$$Af(x) = \sum_{y \sim x} f(y) \quad \text{FOR } f: V(X) \rightarrow \mathbb{C}$$

(REAL SYMMETRIC)

ITS SPECTRUM $\sigma(X) \subset [-3, 3]$

- 3 IS A SIMPLE EIGENVALUE
- -3 IS AN EIGENVALUE IFF X IS BIPARTITE.

$$\sigma(X) : 3 = \lambda_0(X) > \lambda_1(X) \geq \lambda_2(X) \dots \geq \lambda_{n-1}(X) = \lambda_{\min}(X)$$

NB: THE EIGENVALUES ARE TOTALLY REAL ALGEBRAIC INTEGERS!

SPECTRUM OF A ON $\ell^2(T_3)$:

$$\sigma(T_3) = [-2\sqrt{2}, 2\sqrt{2}]$$

KESTEN,
SALLY-SHALIKA
2-ADIC PLANCHEREL
MEASURE.

$$2\sqrt{2} = 2.8284 \dots$$

[4]

WHAT GAPS CAN BE CREATED IN $\sigma(X)$ FOR LARGE X ?

- GAP AT THE TOP (IE 3) IS THE "BASS NOTE" GAP $3 - \lambda_1(X)$ AND IT BEING BOUNDED BELOW BY A POSITIVE CONSTANT IS THE NOTION OF BEING AN EXPANDER.
- TIGHT BINDING HAMILTONIANS IN PHYSICS ASK FOR A GAP AT THE BOTTOM (AT -3).
- IN CHEMISTRY LARGE CARBON CLUSTERS (FULLERENES) THE GAP IN THE MIDDLE IS DECISIVE (HUCKELL ORBITAL STABILITY)




Fig. 6 | The heptagon-kagome device. **a**, Resonator layout (dark blue) and effective lattice (light blue) for a circuit that realizes two shells of the heptagon-kagome lattice. Orange circles indicate three-way capacitive couplers. **b**, Photograph of a physical device that realizes the layout and effective graphs in **a**. The device consists of 140 CPW resonators with fundamental resonance frequencies of 8 GHz, second harmonic frequencies of 16 GHz and a hopping rate of -136.2 MHz at the second harmonic. Four additional CPW lines at each corner of the device couple microwaves into and out of the device for transmission measurements.

Device measurements

We have constructed a device to realize a finite section of the heptagon-kagome lattice. It consists of one central heptagon and two shells of neighbouring tiles, and is shown schematically in Fig. 6a, where each resonator has been approximated by a single line, and the lengths have not been held fixed. The resonators are 7.5 mm long with a fundamental resonance frequency of 8 GHz and a second harmonic of 16 GHz. The second harmonic of this device realizes the heptagon-kagome lattice with a hopping rate of -136.2 MHz. (The fundamental modes of the device obey a different tight-binding model owing to the asymmetry of the mode function within each resonator²². See the Methods for details.)

To minimize parasitic systematic frequency differences between resonator geometries, each resonator type was fabricated individually, and the corresponding resonant frequencies were measured. Commercial microwave simulation packages were unable to achieve the required level of absolute or relative accuracy, so the resonator lengths were then fine-tuned empirically to remove the residual offsets at the level of 30 MHz. For the device shown in Fig. 6b, the average difference between the fundamental frequencies of resonators with different shapes is approximately 0.13% (10 MHz), limited by intrinsic reproducibility within a fabrication run²³ and wire-bonding or parasitic capacitances sensitive to variations between fabrication runs. Each individual shape has a fabrication-induced reproducibility of 0.036% (2.9 MHz), consistent with previous work²³. In addition to the lattice itself, the circuit contains four measurement ports, visible in each corner, which are used to interrogate the lattice.

Theoretical transmission curves for 15 different disorder realizations are shown in Fig. 6c, along with a plot of the experimental transmitted power near the second harmonic frequency of the device. These theoretical curves reproduce most of the qualitative features of the data,

Short stubs protruding inward from the outermost three-way couplers are high-frequency $\lambda/4$ resonators, which maintain a consistent loading of the sites in the outer ring, ensuring uniform on-site energies. **c**, Experimental transmission (S_{21}) for the device in **b** is shown in dark blue. The red curves show theoretical transmission for an ensemble of theoretical models including small systematic offsets in the on-site energies and realistic disorder levels, demonstrating reasonable agreement between theory and experiment.

including the onset of peaks, the location and Fano-like lineshapes of the highest-frequency peaks, and the markedly larger linewidth of the modes near 16.2 GHz which have the largest overlap with the coupling ports. This device therefore demonstrates that hyperbolic lattices can be produced on chip by using CPW resonators, and it paves the way to the study of interactions in hyperbolic space and to simulation of new models with non-constant curvature.

Because of the combination of systematic and random disorder, in practice the flat band will no longer be completely degenerate and will hybridize slightly with the rest of the spectrum. For this heptagon-kagome device, the systematic shape-dependent disorder causes the largest effects: about $0.12|t|$ for the worst shapes and about $0.07|t|$ for typical ones. Random disorder contributes at about the $0.04|t|$ level. Using graph-theoretic studies beyond the scope of the discussion here, we have shown that the bulk gap for the heptagon-kagome lattice is about $0.4|t|$ and that the lower-lying eigenvalues seen in finite-size numerics are whispering-gallery-like edge modes which are very strongly confined to the boundary⁴⁶. Therefore, the gapped flat band of the heptagon-kagome lattice is noticeably broadened, but is able to survive in the experimental realization. These graph-theoretic studies also revealed the existence of closely related and readily realizable lattices with gaps as large as $|t|$ for which the hierarchy of energy scales is favourable.

Conclusion

We have shown that lattices of CPW resonators can be used to produce artificial photonic materials in an effective curved space, including hyperbolic lattices which are typically prohibited as they cannot be isometrically embedded, even in three dimensions. In particular, we conducted numerical tight-binding simulations of hyperbolic analogs of the kagome lattice and demonstrated that they display a flat band

(5)

BASS NOTE SPECTRUM :


FOR $\mathcal{Y} \subset \mathbb{C}$ CUBIC (EG PLANAR)

WE DEFINE THE BASS NOTE SPECTRUM OF \mathcal{Y}

- $BASS_A(\mathcal{Y}) = \{ \lambda_1(x) : x \in \mathcal{Y} \} \subset [-1, 3]$

- IT HAS A DISCRETE PART AND A CONTINUOUS PART.

THEOREM (.)



$BASS_A^D(\text{CUBIC}) = \{ -1, 0, 1, 1, 1, 1, 1.56, \dots \}$ DISCRETE INFINITE IN $[-1, 2\sqrt{2})$

$BASS_A^L(\text{CUBIC}) = [2\sqrt{2}, 3]$

$BASS_A(\text{PLANAR})$ IS RIGID

$BASS_A^D(\text{PLANAR}) = \{ -1, 1, 1, \dots \} \subset [-1, 3]$
DISCRETE

$BASS_A^L(\text{PLANAR}) = \{ 3 \}$

COMMENTS

(5)

(a) FOR PLANAR; $\lambda_1(X) \rightarrow 3$ AS $|X| \rightarrow \infty$
(EG LIPTON-TARJAN)

"PLANAR GRAPHS CANNOT BE EXPANDERS"

(b) FOR $X \in \text{CUBIC}$ J. FRIEDMAN SHOWS THAT

$$\lambda_1(X) \geq 2\sqrt{2} - \frac{100}{(\log X)^2} \Rightarrow \text{BASS (CUBIC)} \text{ IS DISCRETE IN } [-1, 2\sqrt{2}) .$$

(c) SPECIAL RAMANUJAN GRAPHS CONSTRUCTED FROM MODULAR FORMS ENSURE THAT

$$|\text{BASS}_D(\text{CUBIC})| = \infty$$

• N. ZUBRILINA USING WORK OF R. COLEMAN AND B. EDIXHOVEN

SHOWS THAT FOR CERTAIN OF THESE

$$\lambda_1(X_N) \leq 2\sqrt{2} - (1.3)^{-|X(N)|}$$

• MARKUS-SPIELMAN-SRIVASTAVA GIVES X_N 'S WITH

$$\lambda_1(X_N) \leq 2\sqrt{2} - \frac{100}{N}$$

(d) USING THAT THE RANDOM X IN CUBIC IS ALMOST RAMANUJAN (FRIEDMAN) AND THAT ITS EIGENVECTORS ARE DELOCALIZED (H.T. YAU/HUANG...)

F. WEI AND N. ALON SHOW THAT EVERY POINT IN $[2\sqrt{2}, 3]$ IS A LIMIT POINT OF $\lambda_1(X)$ 'S.

GAP AT THE BOTTOM -3 ; HOFFMAN SPECTRUM 14

IF Z IS ANY CONNECTED GRAPH

$L(Z)$ ITS LINE GRAPH:

VERTICES OF $L(Z)$ ARE EDGES OF Z
AND JOIN TWO IF THEY SHARE A VERTEX.

• FACTORIZATION VIA THE INCIDENCE MATRIX \Rightarrow

$$\sigma_A(L(Z)) = \{-2\}^{m-n} \cup \sigma(-2I + A_Z + D_Z)$$

|
DIAGONAL
WITH VALENCE

$\subset [-2, \infty)$

$m = \#$ OF EDGES OF Z

$n = \#$ OF VERTICES

SO $\lambda_{\min}(L(Z)) \geq -2$. "HOFFMAN GRAPH"

FROM $\lambda_{\min}(Z) = \min_{U \neq 0} \frac{\langle U, A_Z U \rangle}{\langle U, U \rangle}$

IT FOLLOWS THAT FOR ANY INDUCED SUBGRAPH B OF Z

$$\lambda_{\min}(Z) \leq \lambda_{\min}(B)$$

SO IF Z IS A HOFFMAN GRAPH THEN (5)
IT CANNOT CONTAIN A HOST OF SMALL
INDUCED MINORS.

\Rightarrow CLASSIFICATION OF HOFFMAN GRAPHS
USING CARTAN MATRICES

CAMERON-GOETHEL-SSEIDEL-SHULT (1975)

"LINE GRAPHS, ROOT SYSTEMS AND ELLIPTIC GEOMETRY"
EXCEPT FOR A FINITE LIST OF SPORADIC
GRAPHS ALL HOFFMAN GRAPHS ARE GENERALIZED
LINE GRAPHS.

• TO CONSTRUCT LINE GRAPHS IN CUBIC
DEFINE $T: \text{CUBIC} \rightarrow \text{CUBIC}$.

FIRST $X \rightarrow S(X)$ BY SUBDIVIDING X ADDING VERTICES
AT THE MIDPOINTS OF EDGES
THIS GIVES A 2-3 REGULAR GRAPH

LET $T(X) := L(S(X)) \in \text{CUBIC}$.

$$|X| T(X) = 3|X|.$$

(EQUIVALENT TO SEWING IN A TRIANGLE AT EACH
VERTEX OF X)

FROM THE CLASSIFICATION OF GRAPHS WITH $\lambda_{\min} \geq -2$ ONE DEDUCES


PROPOSITION (ALICIA KOLLAR, FITZPATRICK, HOUCK, S)

IF $\gamma \in \text{CUBIC}$ AND $\lambda_{\min}(\gamma) \geq -2$ THEN EITHER $\gamma = K_4$ (WHEN $\lambda_{\min}(\gamma) = -1$) OR $\lambda_{\min}(\gamma) = -2$, AND IF γ IS LARGE THEN $\gamma = T(Z)$ FOR SOME $Z \in \text{CUBIC}$.

DEFINE THE HOFFMAN SPECTRUM OF CUBIC GRAPHS TO BE THE VALUES OF λ_{\min} :

$$\text{HOFF}_A(\text{CUBIC}) := \{ \lambda_{\min}(\gamma) : \gamma \in \text{CUBIC} \}$$

IT IS KNOWN (HOFFMAN, ..., YU) THAT



DEFINITIONS: \mathcal{Y} A SUBSET OF CUBIC.

• AN OPEN $U \subset [-3, 3]$ IS A GAP SET FOR \mathcal{Y} IF THERE ARE INFINITELY MANY $X \in \mathcal{Y}$ WITH $\sigma(X) \cap U = \emptyset$.

• A CLOSED $K \subset [-3, 3]$ IS \mathcal{Y} -SPECTRAL IF THERE ARE INFINITELY MANY $X \in \mathcal{Y}$ WITH $\sigma(X) \subset K$.

• $\exists \in [0, 3)$ IS \mathcal{Y} -GAPPED IF \exists HAS A NBH U WHICH IS AN \mathcal{Y} -GAP SET.


THE PREVIOUS PROPOSITION SHOWS THAT $[-3, 2)$ IS A MAXIMAL CUBIC GAP ~~SET~~^{INTERVAL} AND WE SAW THAT $(2\sqrt{2}, 3)$ IS AS WELL.

WE SEEK MAXIMAL GAP SETS OR MINIMAL SPECTRAL SETS AND THEIR DEPENDENCE ON \mathcal{Y} .

SIMILAR QUESTIONS IN ANOTHER SETTING:

- ZEROS OF ZETA FUNCTIONS (OR EIGENVALUES OF FROBENIUS ON COHOMOLOGY) OF CURVES AND ABELIAN VARIETIES A/\mathbb{F}_q OVER A FIXED FINITE FIELD \mathbb{F}_q .
(TFASMAN-VLADUT, DRINFELD, SERRE) IN CONNECTION WITH GOPPA CODES.

$C_q =$
 $|\lambda| = \sqrt{q}$



A-DIMENSION g .
 $2g$ EIGENVALUES
SYMMETRIC
(CONJ INVARIANCE)

SPECTRAL SETS K

- WHAT KIND ~~OF GAPS~~ CAN BE ACHIEVED AS $g \rightarrow \infty$?

FOR CURVES TFASMAN-VLADUT SHOW THAT NO GAPS CAN BE CREATED - RIGID!

FOR ABELIAN VARIETIES A/\mathbb{F}_q , SERRE 2018 SHOWS USING HONDA TATE THEORY THAT THE ONLY CONSTRAINT ON SYMMETRIC SPECTRAL SETS $K \subset C_g$ IS THAT THEIR TRANFINITE DIAMETER OR CAPACITY BE AT LEAST $q^{1/4}$ ($\text{CAP}(C_g) = q^{1/2}$).

• FOR $K \subset \mathbb{C}$ COMPACT ; ITS TRANSFINITE DIAMETER OR CAPACITY IS DEFINED BY

$$n \geq 1 ; \quad d_n(K) = \max_{z_1, \dots, z_n \in K} \prod_{i < j} |z_i - z_j|^{2/(n-1)}$$

$d_n(K)$ IS DECREASING AND $CAP(K) = \lim_{n \rightarrow \infty} d_n(K)$.

THEOREM (FEKETE 1930)

FOR $K \subset \mathbb{C}$ COMPACT ; IF $CAP(K) < 1$

THEN. $\left\{ \alpha : \alpha \text{ AN ALGEBRAIC INTEGER} \right\}$ IS FINITE!
ALL OF WHOSE CONJ
ARE IN K

• RAPHAEL ROBINSON PROVED AN ESSENTIAL CONVERSE FOR SETS $K \subset \mathbb{R}$, THAT IF $CAP(K) \geq 1$ THEN K CONTAINS INFINITELY MANY SUCH TOTALLY REAL ALGEBRAIC INTEGERS!

• SERRE REDUCES THE "WEIL NUMBER" OR EIGEN-VALUES OF FROBENIUS FOR ABELIAN VARIETIES TO ROBINSON'S CONSTRUCTION.

• VERY RECENTLY ALEX SMITH RESOLVED A QUANTITATIVE VERSION OF ROBINSON CONCERNING THE POSSIBLE ^{LIMIT} MEASURES ASSOCIATED TO THE DISTRIBUTION OF THE GALOIS ORBITS CONDENSING ONTO K .

BACK TO SPECTRAL GAPS FOR CUBIC:

9

THEOREM (A. KOLLAR, 5 2021):

(a) ANY CUBIC SPECTRAL SET K
HAS CAPACITY AT LEAST 1.

(b) A CUBIC GAP INTERVAL CAN
HAVE LENGTH AT MOST 2.

(c) EVERY POINT $\lambda \in [-3, 3)$
CAN BE GAPPED WITH PLANAR GRAPHS.

(d). THERE ARE PLANAR CUBIC
SPECTRAL SETS OF CAPACITY 1.

(e) $(-1, 1)$ AND $(-2, 0)$ ARE
MAXIMAL GAP INTERVALS AND
THE FIRST CAN BE GAPPED
WITH PLANAR GRAPHS.

COMMENTS ABOUT PROOFS:

(a) THE LOWER BOUND ON THE CAPACITY OF SPECTRAL SETS HAS ITS ROOTS IN FEKETE.

(b) THE UPPER BOUND^{ON} THE LENGTH OF A GAP INTERVAL IS PROVED COMBINATORIALLY; ONE SHOWS THAT ONE CAN CONSTRUCT AN APPROXIMATE EIGENFUNCTION WITH EIGENVALUE IN A LARGER INTERVAL BY BUILDING ONE IN THE NBH OF A LONG GEODESIC.

(c) THE PROOF THAT THE GAPPABLE SET OF PLANAR GRAPHS IS ALL OF $[-3, 3)$ INVOLVES VARIOUS STEPS:

(i) USING ABELIAN COVERS OVER
(IN FACT SPECIAL ^{LARGE} ACYCLIC COVERS) OF
SMALL MEMBERS OF CUBIC, ONE
ANALYZES INFINITE SUCH TOWERS USING
BLOCH WAVE THEORY (GENERALIZATION
OF FLOQUET THEORY) AND CREAT SOME
GAPS.

(ii) THESE ROOT EXAMPLES ARE
USED TOGETHER WITH THE MAP
 $T: \text{CUBIC} \rightarrow \text{CUBIC}$

TO MOVE THE GAPS AROUND DYNAMICALLY.
THE MOST DIFFICULT REGION TO GAP
IS NEAR 3 SINCE \mathbb{W}^2 ARE PLANAR
GAPPING AND 3 ITSELF CANNOT
BE GAPPED.

THE DYNAMICS ARE USED AS FOLLOWS

THE SPECTRUM OF $T(X)$ IS RELATED TO X (12)

VIA

$$\sigma(T(X)) = f^{-1}(\sigma(X)) \cup \{0\}^{m/2} \cup \{-2\}^{n/2}$$

WHERE

$$f(x) = x^2 - x - 3$$

SO THE DYNAMICS OF f ON \mathbb{R} AND $[-3, 3]$ IS CRITICAL.

$$f^{-1}([-3, 3]) = [-2, 0] \sqcup [1, 3]$$

$$[-3, 3] \supset f^{-1}([-3, 3]) \supset f^{-2}([-3, 3]) \supset \dots$$

$$\Lambda = \bigcap_{m=0}^{\infty} f^{-m}([-3, 3]) \quad \text{IS A CANTOR SET (THE JULIA SET OF } f)$$


$f^m(x) \rightarrow \infty$ AS $m \rightarrow \infty$ IF $x \notin \Lambda$.

$f|_{\Lambda}$ IS TOPOLOGICALLY EQUIVALENT TO THE SHIFT ON $\{0, 1\}^{\mathbb{N}}$.

$$\text{LET } A = \Lambda \cup \bigcup_{m=0}^{\infty} f^{-m}(\{0\})$$

A IS CLOSED AND CONSISTS OF THE CANTOR SET Λ TOGETHER WITH ISOLATED POINTS THAT ACCUMULATE ON Λ .

13



- A IS A MINIMAL SPECTRAL SET,
IT HAS CAPACITY 1 AND
 $\{X \in \text{CUBIC} : \sigma(X) \subset A\}$ CONSISTS OF
FINITELY MANY T -ORBITS (AND X 'S ARE
PLANAR!).

- THE MAXIMAL GAP INTERVALS
 $(-1, 1)$ AND $(-2, 0)$ WERE FOUND
BY ENGINEERING SOME ABELIAN
COVERS AND "FLAT BANDS".

- ANOTHER MINIMAL CUBIC SPECTRAL
SET IS $[-2\sqrt{2}, 2\sqrt{2}] \cup \{3\}$.

THAT THIS SET IS SPECTRAL FOLLOWS
FROM THE EXISTENCE OF RAMANUJAN GRAPHS
THAT IT IS MINIMAL FOLLOWS FROM A THEOREM
OF ABERT-GLASNER-VIRAG
ANY SEQUENCE OF RAMANUJAN GRAPHS
MUST $B-S$ CONVERGE TO T_3 .

\bar{W}_α is contained in $\sigma(\bar{W}_\alpha)$. This follows from G_α being amenable. If Γ_α acts freely on the vertices of \bar{W}_α , i.e. any element $\gamma \neq 1$ in Γ_α fixes none of the vertices of \bar{W}_α , then the quotient $\bar{W}_\alpha/\Gamma_\alpha$ is a multigraph whose spectrum is contained in $\sigma(\bar{W}_\alpha)$. If Γ_α acts without fixing any edges, then the quotient is a graph. We examine each case $\alpha = a, b$ in turn.




FIGURE 13. **Finite planar quotients of \bar{W}_b .** **a**: The infinite graph \bar{W}_b . Four sample involution symmetry points are indicated by black dots. **b**: The quotient obtained with respect to the automorphism σ_O : rotation about O or O' by π . New edges induced by the quotient are indicated in red. In this case, no loops or multiple edges appear. **c**: The quotient with respect to σ_P . In this case, two loops appear. **d**: The quotient with respect to reflection about the central axis. Infinitely many multiple edges appear. **e, f**: The quotient with respect to σ_O and $\sigma_{O'}$, when O and O' are four unit cells apart. This quotient is a planar graph which is $(-1, 1)$ gapped.

Consider first \bar{W}_b . Its automorphism group is generated by four types of elements.

- (i) Translations $t(n)$ by n unit cells. The quotients $\bar{W}_b/\langle t(n) \rangle$ for $n \geq 2$ are the hamburger graphs $W_b(n)$ shown in Fig. 14**b**.
- (ii) The involution σ_O rotating about a central point O by π . Two example points O and O' are shown in Fig. 13**a**. The quotient $\bar{W}_b/\langle \sigma_O \rangle$ is the graph shown in Fig. 13**b**.
- (iii) The involution σ_P rotating about a central point P by π . Two example points P and P' are shown in Fig. 13**a**. The quotient $\bar{W}_b/\langle \sigma_P \rangle$ is a multigraph, shown in Fig. 13**c**.

RIGIDITY:

15

WE RESTRICT TO PLANAR GRAPHS IN CUBIC.

FOR k AN INTEGER LET $\mathcal{F}(k)$ DENOTE THE PLANAR SUCH GRAPHS WITH AT MOST k EDGES PER FACE.

EQUIVALENTLY THEIR DUALS ARE TRIANGULATIONS OF S^2 FOR WHICH THE VERTICES HAVE DEGREE AT MOST k .

• $\mathcal{F}(k)$ IS FINITE FOR $k < 6$ (EULER'S FORMULA)

• $\mathcal{F}(6)$ IS ALREADY QUITE RICH AND CORRESPOND TO WHAT THURSTON CALLS TRIANGULATIONS OF "NON-NEGATIVE CURVATURE". HE PARAMETRIZES THEM IN TERMS OF THE ORBITS OF INTEGER POINTS UNDER THE LINEAR ACTION OF AN ARITHMETIC SUBGROUP OF $SU(9,1)$.

• $\mathcal{F}(k)$, $k \geq 7$ ARE ALREADY VERY RICH.

• THE SUBSET OF $\mathcal{F}(6)$ CONSISTING OF PLANAR CUBIC GRAPHS WITH 6 OR 5 FACES (HEXAGONS AND PENTAGONS - THERE BEING EXACTLY 12 PENTAGONS) ARE CALLED FULLERENES.




FIGURE 2 | A selection of different 3D shapes for regular fullerenes (distribution of the pentagons D_p are set in parentheses). 'Spherically' shaped (icosahedral), for example, (a) $C_{20}-I_h$, (b) $C_{60}-I_h$, and (c) $C_{960}-I_h$ ($D_p = 12 \times 1$); barrel shaped, for example, (d) $C_{140}-D_{3h}$ ($D_p = 6 \times 2$); trigonal pyramidally shaped (tetrahedral structures), for example, (e) $C_{1140}-T_d$ ($D_p = 4 \times 3$); (f) trihedrally shaped $C_{440}-D_3$ ($D_p = 3 \times 4$); (g) nano-cone or menhir $C_{524}-C_1$ ($D_p = 5 + 7 \times 1$); cylindrically shaped (nanotubes), for example, (h) $C_{360}-D_{5h}$, (i) $C_{1152}-D_{6d}$, (j) $C_{840}-D_{5d}$ ($D_p = 2 \times 6$). The fullerenes shown in this figure and throughout the paper have been generated automatically using the *Fullerene* program.³⁵

properties, not least of which is their deep connections to algebraic geometry.¹⁹

Fullerenes have the neat property that the graphs formed by their bond structure are both cubic, planar, and three-connected, for which all faces are either pentagons or hexagons. Because of this, the mathematics describing them is in many cases both rich, simple, and elegant. We are able to derive many properties about their topologies, spatial shapes, surface,

as well as indicators of their chemical behaviors, directly from their graphs.

Planar connected graphs fulfil *Euler's polyhedron formula*,

$$N - E + F = 2 \quad (1)$$

with $N = |\mathcal{V}|$ being the number of vertices (called the *order* of the graph), $E = |\mathcal{E}|$ the number of edges, and $F = |\mathcal{F}|$ the number of faces (for fullerenes these are

THEOREM (ALICIA KOLLAR/FAN WEI/S 2022):

(a) FOR $k \geq 64$ EVERY $\xi \in [-3, 3)$ CAN BE $\mathcal{F}(k)$ GAPPED. WE CONJECTURE THAT THIS CONTINUES TO HOLD FOR $k \geq 7$.

(b) RIGIDITY: THE ONLY POINTS THAT CAN BE $\mathcal{F}(6)$ GAPPED ARE IN $(-1, 1)$ AND THIS INTERVAL IS THE UNIQUE MAXIMAL $\mathcal{F}(6)$ GAP SET.

(c) THE ONLY POINTS THAT CAN BE FULLERENE GAPPED ARE IN

$$J = (-a, b) \cup (b, a)$$

WHERE $a = 0.382\dots$, $b = 0.288\dots$

(a AND b ARE EXPLICIT ALGEBRAIC INTEGERS).

MOREOVER J IS ESSENTIALLY THE UNIQUE MAXIMAL FULLERENE GAP SET.

COMMENTS ON PROOFS

17

• $k \geq 64$. IN ORDER TO LIMIT THE NUMBER OF FACES IN AN ITERATIVE PROCESS OF CONSTRUCTING X 'S IN $\mathcal{Y}(k)$ WITH GAPS (ESPECIALLY NEAR 3) WE SEW IN SOME CAREFULLY CRAFTED ^{SEED} GRAPHS IN THE EDGES OF AN INITIAL GRAPH. THE FORMULAE FOR THE NEW SPECTRA OF THE SEWEN IN GRAPHS INVOLVE RATIONAL FUNCTIONS OF λ AND THEIR ITERATED DYNAMICS ARE STUDIED THROUGH CONTINUED FRACTIONS.

• FOR THE $\mathcal{Y}(6)$ RIGIDITY, WE NEED A DETAILED STUDY OF THE B-S LIMITS OF $\mathcal{Y}(6)$ 'S. THESE CORRESPOND TO INFINITE QUOTIENTS OF THE HEXAGONAL LATTICE, KNOWN AS NANO-TUBES. AN EXPLICIT DETERMINATION OF THEIR SPECTRA AND CONVERGENCE OF SPECTRA.

- FOR FULLERENES THERE IS THE ISSUE OF CAPPING NANO-TUBES WITH PENTAGONS (AND HEXAGONS). THIS LEADS TO THE STUDY OF THE SPECTRA OF INFINITE ONE SIDED NANO-TUBES AND IN PARTICULAR THEIR BOUND STATES.

- THE ^{SPECTRALLY} λ EXTREMAL NANO-TUBE THAT CAN BE FULLERENE CAPPED HAS A UNIQUE ONE SIDED SUCH CAPPING AND THE ^{SINGULAR} λ POINT b IN J THAT CANNOT BE FULLERENE GAPPED CORRESPONDS TO A BOUND STATE.

OPEN QUESTION: WHILE THE THEOREM GIVES A COMPLETE DESCRIPTION OF GAP SETS FOR FULLERENES IT DOES NOT ANSWER THE QUESTION OF WHETHER THE GAP BETWEEN THE TWO MIDDLE EIGENVALUES OF A FULLERENE X \rightarrow THE HOMO-LUMO GAP IN HUCKEL THEORY, MUST CLOSE AS $|X| \rightarrow \infty$? (CARBON CLUSTER STABILITY)

HIGHER RANK S:

VARIOUS RIGIDITY SETS IN . LEADING TO SPECTRAL RIGIDITY .

(1) THE FAMILY \mathcal{Y} OF QUOTIENTS ARE ALL ARITHMETIC (MARGULIS); AND USUALLY EVEN CONGRUENCE SO THE SPECTRA OF ANY \mathbb{P} BECOME PART OF THE GENERAL RAMANUJAN CONJECTURES (DISCUSS IN LECTURE 4).

(2) ANY SEQUENCE OF QUOTIENTS X_j OF \mathcal{S} B-S CONVERGE TO S AS $\text{VOL}(X_j) \rightarrow \infty$

ABERT-BERGERON-BRINGER-GELANDER-NIKOLOV-RAMBAULT-SEMET .

REFERENCES

22

- ABERT-BERGERON-BIRINGER-GELANDER-NIKOLOV-RAIMBAULT-SAMET
ANNALS OF MATH 185 (2017) 2925-2964.
- BOREL-PRASAD PUBL IHES 69 (1989) 119-171
- BUSER-BURGER-DODZIUK LNM 1339, 54-63 (1987)
- BERGER "GEOMETRY REVEALED" SPRINGER (2009).
- BLOMER-BRUMLEY ANN OF MATH 174 (2011) 581-605
- CAMERON-GOETHEL-SSEIDEL-SHULT J. ALGEBRA 43 (1976) 305-327.
- CHAVEL-DODZIUK J. DIFF GEOM 39 (1994) 123-137.
- COLBOIS-COURTOIS ANN. SC. ENS (1991) 507-518.
- COLEMAN-EDIXHOVEN MATH ANN 310 (1998) 119-127.
- FEKETE MATH Z 17 (1923) 228-249
- ELSTRODT-GRUNEWALD-MENNICKE SPRINGER 1998 "GROUPS ACTING ON HYPERBOLIC SPACE."
- FRA CZYK INVENT MATH 224 (2021) 917-985
- FRIEDMAN D.M.J. VOL 69 (1993) 487-525.
- FRIEDMAN PROC 35TH STOC 720-724 (2003)
- HUBER PROC SYMP PURE MATH XXXVI (1979) 181-184.
- HUANG-YAU arXiv 2206.06580 (2022)
- HARTMAN-MAZAC-RASTELLI J. HIGH ENERGY PHYSICS (2019)
- HIDE-MAGEE arXiv 2107.05292 (2021)
- KOLLAR-CARNAK COMM AMER MATH SOC 1 (2021) 1-38.
- KESTEN TAMS 92 (1959) 336-354.
- KRAVCHUK-MAZAC-PAL arXiv 2111.1271 (2021)