




# Assembler la matière quantique atome par atome



**Antoine Browaeys**

*Laboratoire Charles Fabry,  
Institut d'Optique, CNRS, FRANCE*

Collège de France, February 10<sup>th</sup>, 2026



# The program

1. Many-body problem and quantum simulation
2. Example of synthetic quantum system: arrays of individual atoms with Rydberg interactions
3. Examples of simulations: spin models and magnetism

# The context: “many-body problem”


**Goal:** Understand ensembles of **strongly** interacting quantum particles



superfluidity



superconductivity




magnetism



neutron star

Questions

Microscopic description



Difficulty: exponential


**Approximations possible !!**

But... poorly controlled or not valid when interactions dominate

= **Strongly correlated** systems

phase diagram, excitation, thermalization, dynamics, ...

description = models




$$H = \sum_{\langle i,j \rangle, \sigma} t_{ij} c_{j\sigma}^\dagger c_{i\sigma} + U \sum_i n_{i\downarrow} n_{i\uparrow}$$


for  $s = \frac{1}{2}$  :  $N \lesssim 50$

# The context: "many-body problem"


Perovskite:  
solar panels




Catalysis



Batteries &  
cathode



Microscopic description = models




Difficulty: exponential scaling of Hilbert space

**Approximations possible !!**

But... poorly controlled or not valid when interactions dominate

= **Strongly correlated systems**



$$H_{\text{FH}} = -t \sum_{\langle i,j \rangle, \sigma} c_{j\sigma}^\dagger c_{i\sigma} + U \sum_i n_{i\downarrow} n_{i\uparrow}$$


for  $s = \frac{1}{2}$ :  $N \lesssim 50$

# One approach: many-body physics with synthetic quantum systems



R.P. Feynman

Int. J. Theo. Phys. **21** (1982)



Quantum simulation

Georgescu, Rev. Mod. Phys. (2014)

**Well-controlled** quantum systems implementing **many-body Hamiltonians**  
= quantum simulator







Larger tunability than “real” systems (geometry, interactions...)

+

New types of probe & methods (e.g. out-of-equilibrium)

**A new way to look at many-body using quantum information concepts**  
(entanglement...)

# Engineering with individual quantum systems (examples)



Trapped ions

Atoms in optical lattices

Atoms in tweezer arrays

Supercond. Circuits

Photons



IONQ



QUANTINUUM



Pasqal




planqc

IBM Quantum System One


Google AI




# Engineering with individual quantum systems (examples)




Trapped ions




Atoms in optical lattices



Atoms in tweezer arrays



Supercond. Circuits



Photons


**Scalable:** beyond 1000 particles

**Addressability:** local manipulations and measurement

$$\langle \sigma_i^\alpha \rangle, \langle \sigma_i^\alpha \sigma_j^\beta \rangle, \dots$$

**Programmable:** controlled geometry, interactions...

# The team (atom-tweezers-io.org)




**Theory:** E. Demler (ETHZ), A. Bohrdt, F. Grusdt, N. Yao (Harvard), A. Läuchli (Lausanne), T. Roscilde (Lyon), H-P Büchler (Stuttgart), A. Asenjo-Garcia (NY), D. Chang (ICFO), A.-M. Rey (JILA), F. Robicheaux (Perdew)



# Our experiments: many-body physics with synthetic quantum systems

## Tunable arrays of individual Rydberg atoms




$$r \sim 10 \mu\text{m}$$

INSTITUT  
d'OPTIQUE




GRADUATE SCHOOL




$$r \sim \lambda$$

## Light-induced interactions in atomic ensembles




$$r \sim \lambda$$

## Light-induced interactions in arrays of Dy atoms



I. Ferrier-Barbut






PRL 2023, PRA 2024, 2025  
PRXQ 2025

# The program

1. Many-body problem and quantum simulation
2. Example of synthetic quantum system: arrays of individual atoms with Rydberg interactions
3. Examples of simulations: spin models and magnetism

# Assembled arrays of *individual* $^{87}\text{Rb}$ atoms in optical tweezers


Diffraction (spatial light modulator)




Grangier (2001)  
Sortais (2007)  
Bergamini (2004)  
Nogrette (2014)  
Barredo (2016-18)  
Schymik (2021-22)  
Pichard, PR Applied (2024)

# Assembled arrays of *individual* $^{87}\text{Rb}$ atoms in optical tweezers


Diffraction (spatial light modulator)



Fluorescence:  
single shot!!




Grangier (2001)  
Sortais (2007)  
Bergamini (2004)  
Nogrette (2014)  
Barredo (2016-18)  
Schymik (2021-22)  
Pichard, PR Applied (2024)







# Assembled arrays of *individual* $^{87}\text{Rb}$ atoms in optical tweezers

Diffraction (spatial light modulator)




Pasqal



- Grangier (2001)
- Sortais (2007)
- Bergamini (2004)
- Nogrette (2014)
- Barredo (2016-18)
- Schymik (2021-22)
- Pichard, PR Applied (2024)


# Single-atom trapping zoo (2026)



sciencenotes.org

# And now... the interactions

## Arrays of $^{87}\text{Rb}$ atoms




5  $\mu\text{m}$



J. Rydberg  
1854-1919

## Rydberg atoms





Lukin, Zoller... 2000  
Saffman, RMP 2010  
Browaeys, Nat Phys 2020

# And now... the interactions



J. Rydberg  
1854-1919


## Rydberg atoms



Lukin, Zoller... 2000  
Saffman, RMP 2010  
Browaeys, Nat Phys 2020

# And now... the interactions

## Arrays of $^{87}\text{Rb}$ atoms




5  $\mu\text{m}$



J. Rydberg  
1854-1919

## Rydberg atoms



Lifetime  $\propto n^3 > 100 \mu\text{s}$

Transition dipole:  $d_{n, n\pm 1} \sim n^2 e a_0$

$\Rightarrow$  Large dipole-dipole interactions


$$R = 10 \mu\text{m} \Rightarrow V_{\text{int}}/h \sim 1 - 10 \text{ MHz}$$

$\Rightarrow$  timescales  $< \mu\text{sec}$

Lukin, Zoller... 2000  
Saffman, RMP 2010  
Browaeys, Nat Phys 2020

# And now... the interactions

## Arrays of $^{87}\text{Rb}$ atoms




5  $\mu\text{m}$




J. Rydberg  
1854-1919


## Rydberg atoms



## 1<sup>st</sup> demonstrations of controlled Rydberg interactions between two atoms



Nat. Phys. 2009




M. Saffman

Lukin, Zoller... 2000  
Saffman, RMP 2010  
Browaeys, Nat Phys 2020

# And now... the interactions

Arrays of  $^{87}\text{Rb}$  atoms




Rydberg atoms



Rydberg states  $|n, l\rangle$

$$n \gg 1$$




300+ in construction worldwide...

Optical clocks & metrology  
 Quantum simulation  
 Quantum computing  
 Light-matter interface


...

1<sup>st</sup> demonstrations of

... between two atoms



Nat. Phys. 2009





M. Saffman

Lukin, Zoller... 2000  
 Saffman, RMP 2010  
 Browaeys, Nat Phys 2020

# The Pasqal adventure: towards industry grade simulators (2019)

## Engineering



Quantum 2022

## Explore applications

### Quantum Feature Maps for Graph Machine Learning on a Neutral Atom Quantum Processor

Boris Albrecht<sup>1,\*</sup>, Constantin Dalyac<sup>1,2,\*</sup>, Lucas Leclerc<sup>1,3,\*</sup>, Luis Ortiz-Gutiérrez<sup>1,\*</sup>, Slimane Thabet<sup>1,2,\*</sup>, Mauro D'Arcangelo<sup>1</sup>, Vincent E. Elfving<sup>1</sup>, Lucas Lassablière<sup>1</sup>, Henrique Silvério<sup>1</sup>, Bruno Ximenez<sup>1</sup>, Louis-Paul Henry<sup>1</sup>, Adrien Signoles<sup>1</sup>, and Loïc Henriët<sup>1,†</sup>

Phys. Rev. A 2023

### Financial Risk Management on a Neutral Atom Quantum Processor

Lucas Leclerc<sup>1,2,\*</sup>, Luis Ortiz-Gutiérrez<sup>1</sup>, Sebastián Grijalva<sup>1</sup>, Boris Albrecht<sup>1</sup>, Julia R. K. Cline<sup>1</sup>, Vincent E. Elfving<sup>1</sup>, Adrien Signoles<sup>1</sup>, and Loïc Henriët<sup>1,†</sup>

Phys. Rev. Research 2023

### Leveraging Analog Quantum Computing with Neutral Atoms for Solvent Configuration Prediction in Drug Discovery

Mauro D'Arcangelo<sup>1,\*</sup>, Daniele Loco<sup>2,\*</sup>, Fresnel team<sup>1</sup>, Nicolaï Gouraud<sup>2,3,4</sup>, Stanislas Angebault<sup>2</sup>, Jules Sueiro<sup>2</sup>, Pierre Monmarché<sup>3</sup>, Jérôme Forêt<sup>2</sup>, Louis-Paul Henry<sup>1</sup>, Loïc Henriët<sup>1,†</sup>, and Jean-Philip Piquemal<sup>2,4,\*</sup>

Phys. Rev. Research 2024

Dalyac et al. EPJ Quantum Technology (2021) 8:12  
<https://doi.org/10.1146/epjqt.40507-021-00100-3>

EPJ  
ORF



RESEARCH

EPJ Quantum Technology  
a SpringerOpen Journal

Open Access


Qualifying quantum approaches for hard industrial optimization problems. A case study in the field of smart-charging of electric vehicles





# The program

1. Many-body problem and quantum simulation
2. Example of synthetic quantum system: arrays of individual atoms with Rydberg interactions
3. Examples of simulations: spin models and magnetism

# Interactions between Rydberg atoms and spin models





Browaeys & Lahaye, Nat.Phys. (2020)




## Ising model

$$\hat{H} = \sum_{i \neq j} J_{ij} \hat{n}_i \hat{n}_j$$




## Spin 1/2



## XY model



$$\hat{H} = \sum_{i \neq j} J_{ij} (\hat{\sigma}_i^+ \hat{\sigma}_j^- + \hat{\sigma}_i^- \hat{\sigma}_j^+)$$



# Interactions between Rydberg atoms and spin models




Browaeys & Lahaye, Nat.Phys. (2020)



**Spin models = generic systems**  
to study many-body questions:

Quantum phase transition, out-of equilibrium,  
thermalization, topology, entanglement...

$\hat{H} =$



Model

$\hat{\sigma}_j^- + \hat{\sigma}_i^- \hat{\sigma}_j^+$





# From van der Waals interaction to the Ising spin model



van der Waals:  $\Delta E \sim \frac{C_6}{R^6}$

Ex:  $N_2, H_2, O_2$




# From van der Waals interaction to the Ising spin model



van der Waals:  $\Delta E \sim \frac{C_6}{R^6}$

$C_6 \propto n^{11} \Rightarrow$  switchable interaction



Ground state:  $n = 5$   
 Rydberg:  $n = 50$   $\times 10^{11}$

“Equivalent” to interaction between spins




**Ising - like!!**


$$\hat{H}_{\text{int}} = \frac{C_6}{R^6} \hat{n}_1 \hat{n}_2 \sim J \hat{\sigma}_1^z \hat{\sigma}_2^z$$

Rydberg  $n_{1,2} = 1$   
 Ground state  $n_{1,2} = 0$


# From van der Waals interaction to the Ising spin model



# From van der Waals interaction to the Ising spin model



# From van der Waals interaction to the Ising spin model




## Quantum Ising model:

$$H = \frac{\hbar\Omega}{2} \sum_i \hat{\sigma}_x^i + \hbar\delta \sum_i \hat{\sigma}_z^i + \sum_{i < j} \frac{C_6}{R_{ij}^6} \hat{n}_i \hat{n}_j$$

Laser:  $B_{\perp}$   $B_{\parallel}$  spin-spin interactions

# From van der Waals interaction to the Ising spin model



## Quantum Ising model:

$$H = \frac{\hbar\Omega}{2} \sum_i \hat{\sigma}_x^i + \hbar\delta \sum_i \hat{\sigma}_z^i + \sum_{i<j} \frac{C_6}{R_{ij}^6} \hat{n}_i \hat{n}_j$$


Laser:  $B_{\perp}$

$B_{\parallel}$

spin-spin interactions

**Controlled parameters:**  
From negligible to dominant interactions


# An antiferromagnet on a square array



Adiabatic state preparation:

$$H = \lambda H_z + H_{\text{Ising}} \quad \text{Vary } \lambda \gg 1 \rightarrow 0$$

# An antiferromagnet on a square array



$$H = \sum_i \left( \frac{\hbar\Omega(t)}{2} \sigma_x^i - \hbar\delta(t) \hat{n}_i \right) + \sum_{i<j} \frac{C_6}{R_{ij}^6} \hat{n}_i \hat{n}_j$$


Adiabatic state preparation:

$$H = \lambda H_z + H_{\text{Ising}} \quad \text{Vary } \lambda \gg 1 \rightarrow 0$$


**1D:** Pohl PRL 2010; Bloch Science 2015; Lukin Nature 2017, 2019;

**2D:** Lienhard PRX 2018, Bakr PRX 2018; Lukin Nature 2021



# An antiferromagnet on a square array



$$H = \sum_i \left( \frac{\hbar\Omega(t)}{2} \sigma_x^i - \hbar\delta(t) \hat{n}_i \right) + \sum_{i<j} \frac{C_6}{R_{ij}^6} \hat{n}_i \hat{n}_j$$



$\Omega(t), \delta(t)$




$$\frac{C_6}{a^6} \sim \Omega$$




$= |f\rangle$  "bright"



$= |r\rangle$  "dark"



Anti-ferromagnetic ordering!





Ex: MnO, FeO, CoO, NiO, FeCl<sub>2</sub>...



**1D:** Pohl PRL 2010; Bloch Science 2015; Lukin Nature 2017, 2019;  
**2D:** Lienhard PRX 2018, Bakr PRX 2018; Lukin Nature 2021

Scholl et al. Nature (2021)  
 Also: Lukin Nature 2021


# An antiferromagnet on a square array



$\Omega(t), \delta(t)$







**Dynamics: push numerical methods**



State of-the-art simulation (2021):  
MPS limited to 10 x 10  
(14 days on super supercomputer!!)

# Quantum simulation of a real frustrated material: $\text{TmMgGaO}_4$



Theoretical modeling

$$H_{\text{TMGO}} = J_1 \sum_{\langle i,j \rangle} \hat{\sigma}_i^z \hat{\sigma}_j^z + J_2 \sum_{\langle\langle i,j \rangle\rangle} \hat{\sigma}_i^z \hat{\sigma}_j^z + \sum_i \Delta_x \hat{\sigma}_i^x + \Delta_z \hat{\sigma}_i^z,$$





$\Delta_x, \Delta_z$

Liu, PRR 2,043013 (2021)

Simplification

Understand material

# Quantum simulation of a real frustrated material: $\text{TmMgGaO}_4$




## Theoretical modeling

$$H_{\text{TMGO}} = J_1 \sum_{\langle i,j \rangle} \hat{\sigma}_i^z \hat{\sigma}_j^z + J_2 \sum_{\langle\langle i,j \rangle\rangle} \hat{\sigma}_i^z \hat{\sigma}_j^z + \sum_i \Delta_x \hat{\sigma}_i^x + \Delta_z \hat{\sigma}_i^z,$$





$\Delta_x, \Delta_z$

Liu, PRR 2,043013 (2021)



Frustration

# Quantum simulation of a real frustrated material: $\text{TmMgGaO}_4$




## Theoretical modeling

$$H_{\text{TMGO}} = J_1 \sum_{\langle i,j \rangle} \hat{\sigma}_i^z \hat{\sigma}_j^z + J_2 \sum_{\langle\langle i,j \rangle\rangle} \hat{\sigma}_i^z \hat{\sigma}_j^z + \sum_i \Delta_x \hat{\sigma}_i^x + \Delta_z \hat{\sigma}_i^z,$$

Liu, PRR 2,043013 (2021)

## Macroscopic measurement

$$\mu_0 H \sim 14 \text{ T}$$



$$T \sim 20 \text{ mK}$$

$$M(T, H)$$

$$\chi(T, H) = \left( \frac{\partial M}{\partial H} \right)_T$$

$$\Delta_z = \mu_0 H$$

NATIONAL HIGH  
MAGNETIC  
FIELD LABORATORY

Los Alamos  
NATIONAL LABORATORY

## Simulation on Quantum Processor




$$M = \frac{1}{N} \sum_i \langle \hat{\sigma}_i^z \rangle$$

$$\Delta_x = \Omega$$





$$\Delta_z = \delta$$

$$N = 256$$

$$\sim 120 \mu\text{m}$$



# Quantum simulation of a real frustrated material: $\text{TmMgGaO}_4$




## Theoretical modeling

$$H_{\text{TMGO}} = J_1 \sum_{\langle i,j \rangle} \hat{\sigma}_i^z \hat{\sigma}_j^z + J_2 \sum_{\langle\langle i,j \rangle\rangle} \hat{\sigma}_i^z \hat{\sigma}_j^z + \sum_i \Delta_x \hat{\sigma}_i^x + \Delta_z \hat{\sigma}_i^z,$$

Liu, PRR 2,043013 (2021)

## Macroscopic measurement

$$\mu_0 H \sim 14 \text{ T}$$



$$T \sim 20 \text{ mK}$$

$$M(T, H)$$

$$\chi(T, H) = \left( \frac{\partial M}{\partial H} \right)$$


$$H_{\text{TMGO}} \approx 10^{-5} \times H_{\text{QPU}}$$

## Simulation on Quantum Processor




$$N = 256$$

$$\sim 120 \mu\text{m}$$




$$\Delta_z = \delta$$


# Quantum simulation of a real frustrated material: TmMgGaO<sub>4</sub>



Preliminary




# Quantum simulation of a real frustrated material: $\text{TmMgGaO}_4$




Preliminary


## Adiabatic state preparation on QPU




# Quantum simulation of a real frustrated material: $\text{TmMgGaO}_4$



## Adiabatic state preparation on QPU



## Comparison between 3 QPU's & theory





$\text{TmMgGaO}_4$ :  $1/J_1 \sim \text{ps}$

QPU:  $1/J_1 \sim \mu\text{s} \Rightarrow$  slow dynamics

Single-site resolution: magnetic domain



# Resonant interaction between Rydbergs and XY spin model

Browaeys & Lahaye, Nat.Phys. (2020)  
Barredo PRL (2015), de Léséleuc PRL (2017)  
Emperauger PRA (2025)



# Resonant interaction between Rydbergs and XY spin model


Browaeys & Lahaye, Nat.Phys. (2020)  
 Barredo PRL (2015), de Léséleuc PRL (2017)  
 Emperauger PRA (2025)



Non radiative “exchange” of excitation

$$\hat{H}_{XY} = \frac{C_3}{R_{ij}^3} (\hat{\sigma}_i^+ \hat{\sigma}_j^- + \hat{\sigma}_i^- \hat{\sigma}_j^+)$$



$$= \frac{C_3}{2R_{ij}^3} (\hat{\sigma}_i^x \hat{\sigma}_j^x + \hat{\sigma}_i^y \hat{\sigma}_j^y)$$



# Dipolar interactions leads to frustration...


$1/R^3 \Rightarrow$  **frustration!!**

AFM:  $J < 0$






Hard to calculate  
 $N > 100 !!$

Preparation of long-range  
order FM phase (10 x 10)



Chen *et al.*, Nature 2023

Dynamics following quench: spin waves



$1/r^3$  Interaction modifies spin wave dispersion

C. Cheng *et al.*, Science 2025 ; Büchler PRL 2012

# Quantum simulators implementing *spin models* can do more...

## Optimization problems

Cost function = Ising Hamiltonien



$$H = \sum_i h_i n_i + \sum_{i < j} J_{ij} n_i n_j, \quad n_i = 0, 1$$

**Solution** optimization = ground state

**Warning:** competitive w.r. heuristic classical algorithms

$N > 2000 - 5000$  & clock rate  $\sim$  kHz...

arXiv:2502.04291




# Quantum simulators implementing *spin models* can do more...

## Optimization problems

Cost function = Ising Hamiltonien

$$H = \sum_i h_i n_i + \sum_{i < j} J_{ij} n_i n_j, \quad n_i = 0, 1$$


**Solution** optimization = ground state



## Quantum metrology



$^{133}\text{Cs}$



F=4

F=3

1 sec = 9 192 631 770 cycles

States with quantum correlations

$$\frac{\Delta\nu}{\nu_0} \propto \frac{1}{\sqrt{N}} \quad \longrightarrow \quad \frac{\Delta\nu}{\nu_0} \propto \frac{1}{N}$$

Gain x 10 - 100

Article

### Scalable spin squeezing in a dipolar Rydberg atom array

<https://doi.org/10.1038/s41586-023-06414-9>

Received: 14 March 2023

Accepted: 7 July 2023

Guillaume Bornet<sup>1,7</sup>, Gabriel Emperauger<sup>1,7</sup>, Cheng Chen<sup>1,7,8</sup>, Bingtian Ye<sup>2,7</sup>, Maxwell Block<sup>2,7</sup>, Marcus Bintz<sup>2</sup>, Jamie A. Boyd<sup>1</sup>, Daniel Barredo<sup>1,3</sup>, Tommaso Comparin<sup>4</sup>, Fabio Mezzacapo<sup>4</sup>, Tommaso Roscilde<sup>5</sup>, Thierry Lahaye<sup>1</sup>, Norman Y. Yao<sup>2,3,6</sup> & Antoine Browaeys<sup>1</sup>


# Quantum simulators implementing *spin models* can do more...

## Optimization problems

Cost function = Ising Hamiltonien

$$H = \sum_i h_i n_i + \sum_{i < j} J_{ij} n_i n_j, \quad n_i = 0, 1$$


**Solution** optimization = ground state



## Quantum metrology



$^{133}\text{Cs}$




1 sec = 9 192 631 770 cycles

States with quantum correlations

$$\frac{\Delta\nu}{\nu_0} \propto \frac{1}{\sqrt{N}} \quad \longrightarrow \quad \frac{\Delta\nu}{\nu_0} \propto \frac{1}{N}$$

Gain x 10 - 100

**Universal quantum computer**  
(likely long-term...)



- Factoring
- Search algorithm
- ...


# Quantum simulators implementing *spin models* can do more...

## Optimization problems

Cost function = Ising Hamiltonien

$$H = \sum_i h_i n_i + \sum_{i < j} J_{ij} n_i n_j, \quad n_i = 0, 1$$


**Solution** optimization = ground state



## Quantum metrology



$^{133}\text{Cs}$



1 sec = 9 192 631 770 cycles

States with quantum correlations

$$\frac{\Delta\nu}{\nu_0} \propto \frac{1}{\sqrt{N}} \quad \longrightarrow \quad \frac{\Delta\nu}{\nu_0} \propto \frac{1}{N}$$

Gain x 10 - 100

## Universal quantum computer (likely long-term...)

### Article

#### Multi-qubit entanglement and algorithms on a neutral-atom quantum computer

<https://doi.org/10.1038/s41586-022-04603-6>  
 T. M. Graham<sup>1\*</sup>, Y. Song<sup>1\*</sup>, J. Scott<sup>1</sup>, C. Poole<sup>1</sup>, L. Phuttarn<sup>1</sup>, K. Joyn<sup>1</sup>, P. Eichler<sup>1</sup>, X. Jiang<sup>1</sup>, A. Marras<sup>1</sup>, B. Grinkemeyer<sup>2</sup>, M. Kwon<sup>1\*</sup>, M. Ebert<sup>1</sup>, J. Cherek<sup>1</sup>, M. T. Lichtenman<sup>1</sup>, M. Gillette<sup>1</sup>, J. Gilbert<sup>1</sup>, D. Bowman<sup>1</sup>, T. Ballance<sup>1</sup>, C. Campbell<sup>1</sup>, E. D. Dahl<sup>1</sup>, O. Crawford<sup>1</sup>, N. S. Blunt<sup>1</sup>, B. Rogers<sup>1</sup>, T. Noel<sup>1</sup> & M. Saffman<sup>1,2,3\*</sup>  
 Received: 2 January 2022  
 Accepted: 2 March 2022  
 Published online: 20 April 2022


### Article

#### Logical quantum processor based on reconfigurable atom arrays

<https://doi.org/10.1038/s41586-023-06927-3>  
 Dolev Bluvstein<sup>1</sup>, Simon J. Evered<sup>1</sup>, Alexandra A. Geim<sup>1</sup>, Sophie H. Li<sup>1</sup>, Hongyun Zhou<sup>1,2</sup>, Tom Manovitz<sup>1</sup>, Seppele Elaad<sup>1</sup>, Madelyn Cain<sup>1</sup>, Marcin Kalinowski<sup>1</sup>, Dominik Hangleiter<sup>1</sup>, J. Pablo Bonilla Atalides<sup>1</sup>, Nishad Maskara<sup>1</sup>, Iris Cong<sup>1</sup>, Yun Gao<sup>1</sup>, Pedro Sales Rodriguez<sup>1</sup>, Thomas Karolyshyn<sup>1</sup>, Giulia Semeghini<sup>1</sup>, Michael J. Gullans<sup>1</sup>, Markus Greiner<sup>1</sup>, Vladan Vuletić<sup>1</sup> & Mikhail D. Lukin<sup>1,3\*</sup>  
 Received: 21 October 2023  
 Accepted: 1 December 2023  
 Published online: 6 December 2023

# Rydberg based quantum computers today

## Zoning architecture



Single qubit gates

$F > 99.97\%$

Two-qubit gates


$F > 99.2\%$

Proof of principle logical encoding and quantum error detection/correction (2025)

Harvard, Quera, Atom Computing, Inflektion, Pasqal...

Today: circuits with  $\sim 100$  gates

Pasqal




Factorization RSA-2048 (Shor):  $\sim 10^{12}$  gates on  $\sim 10$  million qubits ...

Quantum chemistry ( $\text{Fe}_7\text{MoS}_9\text{Co}$ ): 100-1000 logical qubits  $\sim 10^{14}$  gates



# How large can a *controlled* quantum system be?



??



Zurek, Physics Today 1991



[atom-tweezers-io.org](http://atom-tweezers-io.org)

